

ON THE ORIENTABILITY OF THE SLICE FILTRATION

PABLO PELAEZ

ABSTRACT. Let X be a Noetherian separated scheme of finite Krull dimension. We show that the layers of the slice filtration in the motivic stable homotopy category $\mathcal{SH}$ are strict modules over Voevodsky's algebraic cobordism spectrum. We also show that the zero slice of any commutative ring spectrum in $\mathcal{SH}$ is an oriented ring spectrum in the sense of Morel, and that its associated formal group law is additive. As a consequence, we get that with rational coefficients the slices are in fact motives in the sense of Cisinski-Dégliise [1], and have transfers if the base scheme is excellent. This proves a conjecture of Voevodsky [24, conjecture 11].

2000 *Mathematics Subject Classification.* Primary 14, 55.

Key words and phrases. Algebraic Cobordism, K -theory, Mixed Motives, Oriented Cohomology Theories, Rigid Homotopy Groups, Slice Filtration, Transfers.

1. INTRODUCTION

Let X be a Noetherian separated scheme of finite Krull dimension, and $\mathcal{M}_X$ be the category of pointed simplicial presheaves in the smooth Nisnevich site Sm_X over X equipped with the Quillen model structure [16] introduced by Morel-Voevodsky [7]. We define T in $\mathcal{M}_X$ as the pointed simplicial presheaf represented by $S^1 \wedge \mathbb{G}_m$, where $\mathbb{G}_m$ is the multiplicative group $\mathbb{A}_X^1 - \{0\}$ pointed by 1, and S^1 denotes the simplicial circle. Let $Spt(\mathcal{M}_X)$ denote the category of symmetric T -spectra on $\mathcal{M}_X$ equipped with Jardine's motivic model structure [4]. The homotopy category of $Spt(\mathcal{M}_X)$ is a triangulated category which will be denoted by $\mathcal{SH}$.

Given an integer $q \in \mathbb{Z}$, we consider the following family of symmetric T -spectra

$$C_{eff}^q = \{F_n(S^r \wedge \mathbb{G}_m^s \wedge U_+) \mid n, r, s \geq 0; s - n \geq q; U \in Sm_X\}$$

where F_n is the left adjoint to the n -evaluation functor

$$ev_n : Spt(\mathcal{M}_X) \rightarrow \mathcal{M}_X$$

Voevodsky [24] defines the slice filtration as the following family of triangulated subcategories of $\mathcal{SH}$

$$\dots \subseteq \Sigma_T^{q+1} \mathcal{SH}^{eff} \subseteq \Sigma_T^q \mathcal{SH}^{eff} \subseteq \Sigma_T^{q-1} \mathcal{SH}^{eff} \subseteq \dots$$

where $\Sigma_T^q \mathcal{SH}^{eff}$ is the smallest full triangulated subcategory of $\mathcal{SH}$ which contains C_{eff}^q and is closed under arbitrary coproducts.

It follows from the work of Neeman [8], [9] that the inclusion

$$i_q : \Sigma_T^q \mathcal{SH}^{eff} \rightarrow \mathcal{SH}$$

has a right adjoint $r_q : \mathcal{SH} \rightarrow \Sigma_T^q \mathcal{SH}^{eff}$, and that the following functors

$$f_q : \mathcal{SH} \rightarrow \mathcal{SH}$$

$$s_q : \mathcal{SH} \rightarrow \mathcal{SH}$$

are triangulated, where f_q is defined as the composition $i_q \circ r_q$, and s_q is characterized by the fact that for every $E \in Spt(\mathcal{M}_X)$, we have the following distinguished triangle in $\mathcal{SH}$

$$f_{q+1}E \xrightarrow{\rho_q^E} f_qE \xrightarrow{\pi_q^E} s_qE \longrightarrow \Sigma_T^{1,0} f_{q+1}E$$

We will refer to f_qE as the $(q-1)$ -connective cover of E , and to s_qE as the q -slice of E . It follows directly from the construction that the q -slice of E is right orthogonal with respect to $\Sigma_T^{q+1} \mathcal{SH}^{eff}$, i.e.

$$\mathrm{Hom}_{\mathcal{SH}}(K, s_qE) = 0$$

for every K in $\Sigma_T^{q+1} \mathcal{SH}^{eff}$.

2. STRICT MGL -MODULES

In this section we will show that all the slices have a canonical structure of strict modules in $Spt(\mathcal{M}_X)$ over Voevodsky's algebraic cobordism spectrum.

Let A be a cofibrant ring spectrum with unit in $Spt(\mathcal{M}_X)$, and $A\text{-mod}$ be the category of left A -modules in $Spt(\mathcal{M}_X)$. The work of Jardine [4, proposition 4.19] and Hovey [3, corollary 2.2] implies that the adjunction

$$(A \wedge -, U, \varphi) : Spt(\mathcal{M}_X) \rightarrow A\text{-mod}$$

induces a Quillen model structure $Spt^A(\mathcal{M}_X)$ in $A\text{-mod}$, this means that a map $f : M \rightarrow N$ in $Spt^A(\mathcal{M}_X)$ is a weak equivalence or a fibration if and only if Uf is a weak equivalence or a fibration in $Spt(\mathcal{M}_X)$.

It is easy to see that the homotopy category $\mathcal{SH}^A$ of $Spt^A(\mathcal{M}_X)$ is a triangulated category [13, proposition 3.5.3].

Definition 2.1. *Let E be a spectrum in $\mathcal{SH}$. We say that E is effective if E belongs to the triangulated category $\Sigma_T^0 \mathcal{SH}^{eff}$ defined above.*

Theorem 2.2. *Let A be an effective cofibrant ring spectrum with unit $u^A : \mathbf{1} \rightarrow A$ in $Spt(\mathcal{M}_X)$. If $s_0(u^A)$ is an isomorphism in $\mathcal{SH}$, then for every $q \in \mathbb{Z}$ the functor*

$$s_q : \mathcal{SH} \rightarrow \mathcal{SH}$$

factors (up to a canonical isomorphism) through $\mathcal{SH}^A$

$$\begin{array}{ccc} \mathcal{SH} & \xrightarrow{s_q} & \mathcal{SH} \\ & \searrow \tilde{s}_q & \uparrow UR^A \\ & & \mathcal{SH}^A \end{array}$$

where R^A denotes a fibrant replacement functor in $Spt^A(\mathcal{M}_X)$.

Proof. This follows directly from [13, theorem 3.6.20 and lemma 3.6.21] or [11, theorem 2.1(vi)]. $\square$

The following proposition was proved by Voevodsky [24, p. 10 section 3.4] and Spitzweck [19, corollaries 3.2 and 3.3].

Proposition 2.3 (Voevodsky). *Let MGL denote Voevodsky's algebraic cobordism spectrum [23]. We have that MGL is effective and its unit map $u^{MGL} : \mathbf{1} \rightarrow MGL$ induces an isomorphism on the zero slices in $\mathcal{SH}$*

$$s_0(u^{MGL}) : s_0 \mathbf{1} \xrightarrow{\cong} s_0 MGL$$

Now we can state the main result of this section.

Theorem 2.4. *Let $q \in \mathbb{Z}$ denote an arbitrary integer and E denote an arbitrary symmetric T -spectrum in $Spt(\mathcal{M}_X)$. We have that the q -slice $s_q E$ of E is equipped with a canonical structure of MGL -module in $Spt(\mathcal{M}_X)$. This implies that over any base scheme, the slices are always oriented cohomology theories in the sense of Déglise [2, example 2.12(2)].*

Proof. This follows immediately from theorem 2.2 and proposition 2.3. $\square$

Remark 2.5. *One of the interesting consequences of theorem 2.4 is the fact that over any Noetherian separated base scheme of finite Krull dimension, once we pass to the slices it is possible to apply all the formalism developed by Déglise in [2], e.g. Chern classes and the Gysin triangle.*

3. ORIENTED RING SPECTRA AND FORMAL GROUP LAWS

In this section we will show that given a commutative ring spectrum E in $\mathcal{SH}$, its zero slice s_0E is an oriented ring spectrum (in the sense of Morel [21, definition 3.1]) with additive formal group law in $\mathcal{SH}$. To simplify the notation we will denote by $\mathbb{P}^n$ the trivial projective bundle of rank n over our base scheme X .

Definition 3.1. *Let E be a commutative ring spectrum in $\mathcal{SH}$ with unit*

$$u^E : \mathbf{1} \rightarrow E$$

We say that E is an oriented ring spectrum if there exists an element x_E in $\mathrm{Hom}_{\mathcal{SH}}(F_0(\mathbb{P}^\infty), S^1 \wedge \mathbb{G}_m \wedge E)$, where $\mathbb{P}^\infty$ is the colimit of the diagram

$$\mathbb{P}^1 \rightarrow \mathbb{P}^2 \rightarrow \dots \rightarrow \mathbb{P}^n \rightarrow \dots$$

given by the inclusions of the respective hyperplanes at infinity, such that x_E pulls back to the following composition

$$F_0(\mathbb{P}^1) \cong F_0(S^1 \wedge \mathbb{G}_m) \xrightarrow{id \wedge u^E} F_0(S^1 \wedge \mathbb{G}_m) \wedge E \cong S^1 \wedge \mathbb{G}_m \wedge E$$

The following proposition is classical.

Proposition 3.2 (cf. [14], [15], [21]). *Let (E, x_E) be an oriented ring spectrum in $\mathcal{SH}$, and let $m : \mathbb{P}^\infty \times \mathbb{P}^\infty \rightarrow \mathbb{P}^\infty$ be the map induced by the corresponding Segre embeddings. The pullback of x_E along m , is a formal group law F_E*

$$F_E = \sum_{i+j \geq 1} c_{ij} x^i y^j$$

where the coefficients c_{ij} are elements in the abelian group

$$\mathrm{Hom}_{\mathcal{SH}}(F_0(S^{i+j-1} \wedge \mathbb{G}_m^{i+j-1}), E)$$

and x (resp. y) is the pullback of x_E along the projection in the first factor $p_1 : \mathbb{P}^\infty \times \mathbb{P}^\infty \rightarrow \mathbb{P}^\infty$ (resp. second factor).

Lemma 3.3. *Let E be a commutative ring spectrum in $\mathcal{SH}$ with unit $u^E : \mathbf{1} \rightarrow E$, then its zero slice s_0E is also a commutative ring spectrum in $\mathcal{SH}$, and the induced map*

$$s_0(u^E) : s_0\mathbf{1} \rightarrow s_0E$$

is a map of ring spectra in $\mathcal{SH}$.

Proof. The fact that s_0E is a ring spectrum in $\mathcal{SH}$ follows from [13, theorem 3.6.13], on the other hand the naturality of the pairings constructed in [13, theorem 3.6.9] implies that s_0E is also commutative in $\mathcal{SH}$ and that $s_0(u^E)$ is a map of ring spectra in $\mathcal{SH}$. $\square$

Lemma 3.4. *The natural map*

$$\pi_0^{MGL} : MGL \cong f_0MGL \rightarrow s_0MGL$$

is a map of ring spectra in $\mathcal{SH}$.

Proof. By proposition 2.3 we have that MGL is naturally isomorphic to f_0MGL in $\mathcal{SH}$. On the other hand, theorem 3.6.10(3) in [13] implies that π_0^{MGL} is a map of ring spectra in $\mathcal{SH}$. $\square$

Theorem 3.5. *Let E be a commutative ring spectrum in $\mathcal{SH}$. Then its zero slice s_0E is an oriented ring spectrum in the sense of Morel and it is equipped with a canonical orientation given by the following composition*

$$MGL \xrightarrow{\pi_0^{MGL}} s_0MGL \xrightarrow{(s_0(u^{MGL}))^{-1}} s_0\mathbf{1} \xrightarrow{s_0(u^E)} s_0E$$

Furthermore, the associated formal group law F_{s_0E} of s_0E is additive.

Proof. The universality of MGL (cf. [21, theorem 4.3], [10, theorem 2.7] and [20, proposition A.2]) implies that in order to show that the map defined above gives an orientation for s_0E , it is enough to see that all the maps are in fact maps of ring spectra in $\mathcal{SH}$. But this follows directly from proposition 2.3 together with lemmas 3.3 and 3.4.

On the other hand, the formal group law of s_0E

$$F_{s_0E} = \sum_{i+j \geq 1} c_{ij} x^i y^j$$

has coefficients c_{ij} which by construction are in the abelian group

$$\mathrm{Hom}_{\mathcal{SH}}(F_0(S^{i+j-1} \wedge \mathbb{G}_m^{i+j-1}), s_0E)$$

However, if $i+j > 1$ then $F_0(S^{i+j-1} \wedge \mathbb{G}_m^{i+j-1})$ is automatically in $\Sigma_T^1 \mathcal{SH}^{eff}$; hence

$$\mathrm{Hom}_{\mathcal{SH}}(F_0(S^{i+j-1} \wedge \mathbb{G}_m^{i+j-1}), s_0E) = 0$$

since s_0E is right orthogonal with respect to $\Sigma_T^1 \mathcal{SH}^{eff}$.

Therefore, the formal group law of s_0E

$$F_{s_0E} = \sum_{i+j \geq 1} c_{ij} x^i y^j = x + y$$

is additive, as we wanted. □

4. APPLICATIONS

In this section we will show that with rational coefficients all the slices $s_q(E) \otimes \mathbf{1}_{\mathbb{Q}}$ are in a natural way motives in the sense of Cisinski-Déglise [1]; as a consequence we will get that over an excellent base scheme the presheaves of rational rigid homotopy groups have transfers, this proves a conjecture of Voevodsky [24, conjecture 11].

4.1. Slices and the Cisinski-Déglise category of motives. Let

$$\mathbf{H}_{\mathbb{B},X} = KGL_X^{(0)} \in Spt(\mathcal{M}_X)$$

denote the Beilinson motivic cohomology spectrum constructed by Riou in [17]. The work of Cisinski-Déglise shows in particular that $\mathbf{H}_{\mathbb{B},X}$ is a commutative cofibrant ring spectrum in $Spt(\mathcal{M}_X)$ (cf. [1, corollary 13.2.6]); and that the homotopy category of $\mathbf{H}_{\mathbb{B},X}$ -modules $\mathcal{SH}^{\mathbf{H}_{\mathbb{B},X}}$ is naturally equivalent to the Cisinski-Déglise category of motives $DM_{\mathbb{B},X}$ (cf. [1, theorem 13.2.9]).

Theorem 4.1. *If we consider rational coefficients, the zero slice of the sphere spectrum $s_0(\mathbf{1}) \otimes \mathbf{1}_{\mathbb{Q}}$ is equipped with a unique structure of $\mathbf{H}_{\mathbb{B},X}$ -algebra in $Spt(\mathcal{M}_X)$.*

In particular, there exists a unique map $\eta_{s_0(\mathbf{1}) \otimes \mathbf{1}_{\mathbb{Q}}}$ of ring spectra in $\mathcal{SH}$ such that the following diagram is commutative

$$\begin{array}{ccccc} MGL \otimes \mathbf{1}_{\mathbb{Q}} & \xrightarrow{\pi_0^{MGL} \otimes id} & s_0(MGL) \otimes \mathbf{1}_{\mathbb{Q}} & \xrightarrow{(s_0(u^{MGL}))^{-1} \otimes id} & s_0(\mathbf{1}) \otimes \mathbf{1}_{\mathbb{Q}} \\ \eta_H \downarrow & & \nearrow \eta_{s_0(\mathbf{1}) \otimes \mathbf{1}_{\mathbb{Q}}} & & \\ \mathbf{H}_{\mathbb{B},X} & & & & \end{array}$$

Proof. By theorem 3.5 we have that $s_0(\mathbf{1})$ is orientable. Therefore, the result follows directly from corollary 13.2.15(Ri),(Rii),(Riii) in [1]. $\square$

Theorem 4.2. *Let $q \in \mathbb{Z}$ denote an arbitrary integer and E denote an arbitrary symmetric T -spectrum in $Spt(\mathcal{M}_X)$. We have that the q -slice of E with rational coefficients $s_q(E) \otimes \mathbf{1}_{\mathbb{Q}}$ is equipped with a canonical structure of $\mathbf{H}_{\mathbb{B},X}$ -module in $Spt(\mathcal{M}_X)$. This implies that over any base scheme, the slices with rational coefficients $s_q(E) \otimes \mathbf{1}_{\mathbb{Q}}$ are motives in the sense of Cisinski-Déglise.*

Proof. Using corollary 13.2.15(i),(iv),(v) in [1], we get that it is enough to show that $s_q(E) \otimes \mathbf{1}_{\mathbb{Q}}$ is a $\mathbf{H}_{\mathbb{B},X}$ -module in $\mathcal{SH}$. On the other hand, by theorem 4.1 we just need to check that $s_q(E) \otimes \mathbf{1}_{\mathbb{Q}}$ is a $s_0(\mathbf{1}) \otimes \mathbf{1}_{\mathbb{Q}}$ -module in $\mathcal{SH}$. Finally, this follows directly from theorem 3.6.14(6) in [13]. $\square$

4.2. Rational Rigid Homotopy groups. Given a symmetric T -spectrum E in $Spt(\mathcal{M}_X)$, Voevodsky defines the presheaves of rigid homotopy groups $\pi_{p,q}^{rig}(E)$ on Sm_X as follows:

$$\begin{aligned} \pi_{p,q}^{rig}(E) : Sm_X &\longrightarrow \text{Abelian Groups} \\ U &\longmapsto \text{Hom}_{\mathcal{SH}}(F_0(S^p \wedge \mathbb{G}_m^q \wedge U_+), s_q E) \end{aligned}$$

Conjecture 11 in [24] claims that these presheaves have transfers.

Theorem 4.3. *Let $p, q \in \mathbb{Z}$ denote arbitrary integers and E denote an arbitrary symmetric T -spectrum in $Spt(\mathcal{M}_X)$. Furthermore, assume that the base scheme*

X is excellent. Then the presheaves of rigid homotopy groups of E with rational coefficients $\pi_{p,q}^{rig}(E) \otimes \mathbb{Q}$ have transfers.

Proof. Clearly, it suffices to show that $s_q(E) \otimes \mathbf{1}_{\mathbb{Q}}$ has transfers. Now, theorem 4.2 implies that $s_q(E) \otimes \mathbf{1}_{\mathbb{Q}}$ is in $DM_{\mathbb{B},X}$. Since we are assuming that X is excellent, theorem 15.1.2 in [1] implies that $DM_{\mathbb{B},X}$ is naturally equivalent to the category of motives $DM_{qfh,X}$ constructed using qfh -sheaves with rational coefficients. Thus, the result follows from theorem 3.3.8 in [22] (see also proposition 9.5.5 in [1]) which implies that every qfh -sheaf is canonically equipped with transfers. $\square$

Remark 4.4. *Theorem 4.3 was proved using the functoriality of the slice filtration in [12, theorem 4.4] for schemes defined over a field of characteristic zero; on the other hand, if the base scheme X is smooth over a perfect field k , then theorem 4.3 holds even with integral coefficients (cf. [13, theorem 3.6.22]). Both proofs rely on the computation of Levine [5] and Voevodsky [25] for the zero slice of the sphere spectrum, as well as on the work of R ndigs- stvær [18].*

The analogue of this question for the category of S^1 -spectra is studied by Levine in [6].

ACKNOWLEDGEMENTS

The author would like to warmly thank Frédéric Déglise for several useful conversations and suggestions, as well as for putting in our hands the technical tools from [2] and [1]; and also thank Denis-Charles Cisinski for bringing to our attention the argument which allowed us to extend theorem 4.3 from geometrically unibranch base schemes to arbitrary excellent schemes.

REFERENCES

- [1] D.-C. Cisinski and F. Déglise. Triangulated categories of mixed motives. *preprint*, 2009.
- [2] F. Déglise. Around the Gysin triangle. II. *Doc. Math.*, 13:613–675, 2008.
- [3] M. Hovey. Monoidal model categories. *preprint*, 1998.
- [4] J. F. Jardine. Motivic symmetric spectra. *Doc. Math.*, 5:445–553 (electronic), 2000.
- [5] M. Levine. The homotopy coniveau tower. *J. Topol.*, 1(1):217–267, 2008.
- [6] M. Levine. Slices and transfers. *Doc. Math.*, Extra volume: Andrei A. Suslin’s Sixtieth Birthday:393–443, 2010.
- [7] F. Morel and V. Voevodsky. $\mathbf{A}^1$ -homotopy theory of schemes. *Inst. Hautes Études Sci. Publ. Math.*, (90):45–143 (2001), 1999.
- [8] A. Neeman. The Grothendieck duality theorem via Bousfield’s techniques and Brown representability. *J. Amer. Math. Soc.*, 9(1):205–236, 1996.
- [9] A. Neeman. *Triangulated categories*, volume 148 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2001.
- [10] I. Panin, K. Pimenov, and O. Röndigs. A universality theorem for Voevodsky’s algebraic cobordism spectrum. *Homology, Homotopy Appl.*, 10(2):211–226, 2008.
- [11] P. Pelaez. Mixed motives and the slice filtration. *C. R. Math. Acad. Sci. Paris*, 347(9-10):541–544, 2009.
- [12] P. Pelaez. On the functoriality of the slice filtration. *preprint*, 2010.
- [13] P. Pelaez. Multiplicative properties of the slice filtration. *Astérisque*, (335):xvi+291 pp., 2011.
- [14] D. Quillen. On the formal group laws of unoriented and complex cobordism theory. *Bull. Amer. Math. Soc.*, 75:1293–1298, 1969.
- [15] D. Quillen. Elementary proofs of some results of cobordism theory using Steenrod operations. *Advances in Math.*, 7:29–56 (1971), 1971.
- [16] D. G. Quillen. *Homotopical algebra*. Lecture Notes in Mathematics, No. 43. Springer-Verlag, Berlin, 1967.
- [17] J. Riou. Algebraic K -theory, $\mathbf{A}^1$ -homotopy and Riemann-Roch theorems. *J. Topol.*, 3(2):229–264, 2010.
- [18] O. Röndigs and P. A. Østvær. Modules over motivic cohomology. *Adv. Math.*, 219(2):689–727, 2008.
- [19] M. Spitzweck. Relations between slices and quotients of the algebraic cobordism spectrum. *Homology, Homotopy Appl.*, 12(2):335–351, 2010.
- [20] M. Spitzweck and P. A. Østvær. The Bott inverted infinite projective space is homotopy algebraic K -theory. *Bull. Lond. Math. Soc.*, 41(2):281–292, 2009.
- [21] G. Vezzosi. Brown-Peterson spectra in stable $\mathbf{A}^1$ -homotopy theory. *Rend. Sem. Mat. Univ. Padova*, 106:47–64, 2001.
- [22] V. Voevodsky. Homology of schemes. *Selecta Math. (N.S.)*, 2(1):111–153, 1996.
- [23] V. Voevodsky. $\mathbf{A}^1$ -homotopy theory. In *Proceedings of the International Congress of Mathematicians, Vol. I (Berlin, 1998)*, number Extra Vol. I, pages 579–604 (electronic), 1998.
- [24] V. Voevodsky. Open problems in the motivic stable homotopy theory. I. In *Motives, polylogarithms and Hodge theory, Part I (Irvine, CA, 1998)*, volume 3 of *Int. Press Lect. Ser.*, pages 3–34. Int. Press, Somerville, MA, 2002.
- [25] V. Voevodsky. On the zero slice of the sphere spectrum. *Tr. Mat. Inst. Steklova*, 246(Algebr. Geom. Metody, Svyazi i Prilozh.):106–115, 2004.

UNIVERSITÄT DUISBURG-ESSEN, MATHEMATIK, 45117 ESSEN, GERMANY
E-mail address: pablo.pelaez@uni-due.de